



Antioxidant and Antibacterial Activities of Curd and Whey Kefir Produced from Etawa Goat Milk

Siwi Meutia Sadewi, Nurhasanah Nurhasanah*, Sudibyo Sudibyo, Neneng Windayani, Agung Abadi Kiswandono, and Heri Satria

Received : August 21, 2023

Revised : December 1, 2023

Accepted : December 6, 2023

Online : December 11, 2023

Abstract

Kefir is a fermented drink with many benefits. Kefir contains antioxidant compounds that can inhibit free radical activity. In addition, kefir contains lactic acid bacteria, which provide good antibacterial activity and improve the health of the digestive tract. This research aimed to determine the antioxidant and antibacterial activities of kefir made from Etawa goat milk. The methods used in this research included fermentation of Etawa goat milk with 10 % kefir seeds at 37 °C and a fermentation time of 2 days. Antioxidant activity was tested using 1,1-diphenyl-2-picrylhydrazyl (DPPH), and antibacterial activity was tested using agar diffusion. The peptide molecular weights in kefir were analyzed using SDS-PAGE. The results showed that the antibacterial and antioxidant activities in Etawa goat's milk curd kefir has a higher activity than whey kefir. The antioxidant activity of curd has an IC₅₀ of 43.99 ppm, which is a powerful antioxidant. Curd's antibacterial activity forms a clear zone of 7 mm against *Bacillus cereus* and is included in the solid antibacterial category. In *Escherichia coli*, a clear zone of 6 mm was formed and included in the moderate antibacterial category. The results of SDS-PAGE gel visualization found three prominent bands measuring 10, 15, and 33 kDa, respectively.

Keywords: etawa goat milk, curd and whey, antioxidant, antibacterial

1. INTRODUCTION

Kefir is a fermented milk product that is popular in many countries because kefir is a probiotic drink that contains many functional substances that are good for health [1]. Kefir differs from other fermented kinds of milk in that it is formed from the fermentation process of kefir grains, which contain a mixture of lactic acid bacteria and yeast. In addition, kefir seeds also contain lactic acid bacteria such as *Lactobacillus brevis*, *L. helveticus*, *L. kefir*, *Leuconostoc mesenteroides*, *Kluyveromyces lactis*, and *K. marxianus* [2]. The microorganisms that make up the kefir seeds have antibacterial activity, which can assist the digestive system and inhibit pathogenic microorganisms [3]. Kefir has a therapeutic effect because it contains bioactive components and can potentially maintain body health, such as by helping prevent colon cancer and lowering cholesterol levels in the blood. It also has antibacterial properties and can ward off free

radicals by functioning as an antioxidant [4].

Free radicals can form inside the body or come from outside the body. Free radicals are formed in the body due to metabolic processes, such as digesting food and using oxygen. Free radicals may come from smoke exposure to radiation, toxic substances, and heavy metals (pollution) [5][6]. Sources of antioxidants are needed to treat cells exposed to free radicals. The exploration of natural sources of antioxidants continues with the growing human awareness of the dangers of free radicals [7]. Therefore, there is increasing worldwide interest in finding natural, food-based antioxidants capable of protecting the human body from free radical attack and inhibiting the development of many chronic diseases [8].

Kefir can be made from various raw materials, such as cow, soy, and goat. During kefir processing, two products are formed, whey and curd [9]. In addition, Etawa goat milk is hydrolyzed with enzymes or by fermentation containing bioactive peptides. Bioactive peptides are a collection of 2–50 amino acids bound to each other by peptide bonds, which have health benefits. Bioactive peptides can be released from their parent proteins through hydrolysis by proteolytic enzymes or fermentation processes [10]. Bioactive peptides produced during fermentation have antihypertensive, antibacterial, antioxidant, antimicrobial, and anti-inflammatory activities. Bioactive peptides also have good functions for body health, such as biological activity in the digestive, immune, and nervous

Publisher's Note:

Pandawa Institute stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright:

© 2023 by the author(s).

Licensee Pandawa Institute, Metro, Indonesia. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).




Figure 1. Etawa goat milk kefir (a) whey and curd before being separated; (b) whey and (c) curd.

systems [11].

Producing kefir using sheep's milk as a raw material has obtained antioxidants ranging from 18.39 to 72.41% [12]. Apart from that, it is also known that goat's milk kefir has antibacterial properties against *Propionibacterium acnes* bacteria, forming an inhibition zone between 8.17 to 9.50 mm [13]. Other research reports that goat's milk kefir shows inhibitory activity against *Staphylococcus aureus* bacteria [14], and it is known that goat's milk kefir has antibacterial potential against the pathogens *E. coli* ATCC 8739 and *Salmonella enteric subsp. enterica serovar Typhimurium* ATCC 14028 [15]. Based on the background of the description above, the antioxidant and antibacterial activity of Etawa goat's milk kefir fermentation products and the molecular weight of kefir peptides were identified in this research.

2. MATERIALS AND METHODS

2.1. Materials

Etawa goat milk taken from the Telaga Risqi farm East Metro Lampung, kefir grains (UIN Sunan Gunung Djati Bandung), 1,1-diphenyl-2-picrylhydrazyl (DPPH), NaCl (Merck), *B. cereus*

and *E. coli* bacteria cultures (Laboratorium of Microbiology Lampung University), Nutrient Agar media (Merck), and chloramphenicol (Forma mart) were used in this work.

2.2. Methods

2.2.1. Etawa Goat Milk Kefir Production

Etawa goat milk kefir is made by pasteurizing it at 72 °C for 15 s, then leaving it at room temperature until it reaches 30 °C [16]. Etawa goat milk pasteurized is taken in as much as 1 L and added with 10% kefir seeds. The kefir milk is fermented at 37 °C for 48 h [17].

2.2.2. Antioxidant Test

The antioxidant test was carried out using the DPPH method. 5 mL of the sample was dissolved in 5 mL of methanol, which was used as the mother liquor. For a 100 ppm primary solution, a series of samples were made with 100, 80, 60, 40, and 20 ppm concentrations. A sample solution with each concentration was put into a test tube, and 2 mL of 40 ppm DPPH was added. The sample solution was vortexed until homogeneous and incubated for 30 minutes in a dark room. The color change from purple to yellow was then observed. Then, the absorbance value of the sample solution was measured with a UV-Vis spectrophotometer at a wavelength of 512–520 nm. The blank solution was 2 mL of methanol and 2 mL of 40 ppm DPPH [18].

2.2.3. Antibacterial Test

The antibacterial test was carried out using the agar diffusion method. A suspension of *E. coli* and

Table 1. IC₅₀ value of Etawa goat milk kefir.

Sample	IC ₅₀ (ppm)
Ascorbic acid (control)	5.443
Whey	46.01
Curd	43.99

B. cereus bacteria was put into 0.9 mL of NaCl, then vortexed and equalized to the McFarland standard. The agar media that had been prepared was added to 0.1 mL of the bacterial suspension and then levelled with a swab. Then, put the disc dripped with samples and a positive control on it and incubate it for 24 h at 37 °C. The positive control used was chloramphenicol at 2.5 mg/mL. The clear zone around the paper disc indicates a positive test. The clear zone formed was measured and compared with the antibiotic chloramphenicol [4].

2.2.4. Peptide Weight Analysis

SDS-PAGE electrophoresis can be used to determine the molecular weight of the bioactive peptides being analyzed [19]. SDS PAGE electrophoresis (sodium dodecyl sulfate-polyacrylamide gel electrophoresis) The fermented protein samples were denatured with buffer (1 M Tris-HCl pH 6.8, 20% SDS, NO₃) with a protein-buffer ratio of 2:1, boiled at 90 °C for 10 min, and centrifuged for 5 min. An electrophoresis apparatus was prepared. Polyacrylamide gel was prepared from a solution of acrylamide (48%), bisacrylamide (1.5%), stacking buffer (Tris-HCl 0.5 M pH 6.8), resolving buffer (Tris-HCl 1.5 M pH 8.8), 10% SDS, APS, and TEMED as catalysts. After the bottom gel (resolving gel) is formed, stacking gel is inserted at the top, and a mold is made to place the sample protein. Sample electrophoresis was carried out at 150 V for 60 min. The staining protein is Coomassie Blue. The staining results are washed in a destaining solution [20].

3. RESULTS AND DISCUSSIONS

3.1. Etawa Goat's Milk Kefir

Etawa goat milk kefir is a probiotic drink that has many health benefits. In this research,

processed milk kefir products were made using Etawa goat milk as raw material. The results obtained from making Etawa goat milk kefir with 10 % kefir seeds for 48 h at 37 °C form two parts, i.e., whey and curd. The results of making Etawa goat milk kefir can be seen in Figure 1.

Figure 1 shows that after 48 h of fermentation, Etawa goat milk kefir forms two parts: the lower layer (the clear phase) is called whey, and the top layer (the solid phase) is called curd. The stage of making kefir goes through a fermentation process to produce a solid called curd and a transparent layer in the form of a liquid called whey [21].

3.2. Antioxidant Activity

The antioxidant activity test in this research used the DPPH method, indicated in IC₅₀ values, to ascertain the antioxidant capacity of whey and curd kefir in Etawa goat milk. The antioxidant activity can be divided into five classes based on the IC₅₀ value, according to Molyneux [22]: very strong (IC₅₀ < 50 ppm), strong (50 < IC₅₀ < 100 ppm), moderate (100 < IC₅₀ < 150 ppm), weak (150 < IC₅₀ < 200 ppm), and very weak (IC₅₀ > 200 ppm). Whey and curd in this research showed IC₅₀ values ranging from 43.99 to 46.01 ppm, which included very potent antioxidants. Table 1 displays the findings of antioxidant testing on fermented milk products made from goat whey and curd from Etawa.

The results in the table show that curd has higher antioxidant activity because it has a higher total lactic acid value than whey. In this research, whey and curd had IC₅₀ values between 43.99–46.01 ppm, including powerful antioxidants. High total lactic acid can increase antioxidant activity due to the formation of organic acids produced by lactic acid bacteria and yeast, which are synergistic by providing H⁺ ions to free radicals [23]. Fermented milk contains natural antioxidants, which are useful

Table 2. Diameter of clear zone whey and curd kefir against *Bacillus cereus* bacteria.

Sample	Clear Zone Diameter (mm)			Average (mm)
	I	II	III	
Whey	4	4	4	4
Curd	7	8	6	7
Chloramphenicol	24	24	24	24

Table 3. Diameter of clear zone whey and curd kefir against *Escherichia coli* bacteria.

Sample	Clear Zone Diameter (mm)			Average (mm)
	I	II	III	
Whey	3	3	3	3
Curd	7	6	5	6
Chloramphenicol	24	24	24	24

for human health because they are safer and more easily absorbed by the body than synthetic antioxidants [7]. Another research used various kinds of milk, such as cows and donkeys. It was found that all fermented kefir products containing amino acids from the breakdown of milk protein by lactic acid bacteria can produce antioxidant activity that is good for health [24].

3.3. Antibacterial Activity

This study examined the whey and curd kefir's antimicrobial properties in Etawa goat milk. At this point, the antibacterial test aims to ascertain whether whey and curd can inhibit both Gram-positive and Gram-negative bacteria. The agar diffusion method was used to conduct the antibacterial test. *E. coli* was employed for Gram-negative bacteria and *B. cereus* for Gram-positive bacteria in the bacterial solution. A clear zone forming around the disc is a sign of antibacterial action. The formed clear zone denotes the region where bacterial growth is inhibited. The results of testing the antibacterial activity of whey and curd kefir in Etawa goat milk with a suspension of *B. cereus* and *E. coli* bacteria can be seen in Tables 2 and 3.

Based on the data in Table 2, the diameter of the clear zone between whey and kefir curd against *B. cereus* bacteria is 4 and 7 mm. As for Table 3, the diameter of the clear zone between whey and kefir curd against *E. coli* bacteria is 3 and 6 mm. Based on these results, it can be seen that the antibacterial activity of whey and kefir curd against *E. coli* was weaker compared to *B. cereus* bacteria, characterized by the formation of a smaller clear zone diameter. This is because *E. coli* is a group of Gram-negative bacteria that have more complex walls. The cell wall of Gram-negative bacteria has as much as 10 % peptidoglycan and several peptide cross-links. The outer layer consists of proteins,

lipoproteins, and lipopolysaccharides. This complex *E. coli* cell wall makes it more difficult for antibacterial compounds to penetrate [25]. The ability of goat's milk kefir to have antibacterial activity using 5% seeds gave clear zone results of 2.19 mm against *E. coli* bacteria and 2.50 mm against *S. aureus* bacteria [13].

The inhibition zone, which is antibacterial activity, is categorized into three criteria. The first criterion is an inhibition zone with a weak category with a zone area of 0–3 mm; an obstacle zone with a moderate category with a zone area of 3–6 mm; and the third criterion is an inhibition zone with a strong category with an area of inhibition above 6 mm [26]. Based on the results obtained in this research, it can be concluded that the antibacterial activity of curd forms a clear zone of 7 mm against *B. cereus* and is included in the solid antibacterial category. In *E. coli*, a clear zone of 6 mm was formed and included in the moderate antibacterial category.

3.4. Whey and Curd Kefir Bioactive Peptide Weights

Determination of the peptide molecular weight




Figure 2. Etawa goat milk kefir peptide weight (M) spectra multicolor broad range protein ladder 26634 for (A) whey; and (B) curd.

was carried out on the whey and curd kefir fractions using the SDS-PAGE method. The results of the analysis of the molecular weight of Etawa goat's milk kefir peptides using SDS PAGE are presented in Figure 2.

Figure 2 shows that the Etawa goat's milk kefir peptide weight intensity in whey and curd has three prominent bands measuring 10, 15, and 33 kDa. Whereas in Etawa goat's milk curd, one other band has a reasonably high peptide intensity, namely the 24 kDa band. This shows that Etawa goat's milk kefir curd has more varied peptides and that the curd has more molecular weight variations. The weight of other peptides with very low intensity in whey and curd of Etawa's milk appears in the 50–100 kDa range. The peptide weighs 10–38 kDa in fermented goat milk yoghurt and has constituent amino acids such as valine, lysine, glutamate, methionine, and alanine [27]. The peptide weight between 10–70 kDa belongs to the water-soluble protein group and is generally a glycoprotein group obtained from protein hydrolysis in goat's milk [20], while the band detected at 50–80 kDa weight belongs to the lactoferrin group obtained from goat's milk kefir Etawa [28].

4. CONCLUSIONS

Antibacterial and antioxidant activities in Etawa goat's milk curd kefir is higher than whey kefir. Curd's antibacterial activity against *Bacillus cereus* was included in the category of intense antibacterial activity while the antibacterial activity against *Escherichia coli* was included in the moderate antibacterial category. The antioxidant activity of curd has an IC₅₀ of 43.99 ppm, categorized as a very strong antioxidant. The molecular weight of kefir peptides in whey and curd has three prominent bands, namely 10, 15, and 33 kDa. In curd kefir, there is one band weighing 24 kDa, which has a reasonably high intensity.

AUTHOR INFORMATION

Corresponding Author

Nurhasanah Nurhasanah — Department Chemistry, Lampung University, Bandar Lampung-35141 (Indonesia);

 orcid.org/0000-0002-5552-7665

Email: nur.hasanah@fmipa.unila.ac.id

Authors

Siwi Meutia Sadewi — Magister of Chemistry Program, Lampung University, Bandar Lampung -35141 (Indonesia);

 orcid.org/0000-0002-5525-5073

Sudibyo Sudibyo — Research Center for Mining Technology, National Research and Innovation Agency (BRIN), South Lampung-35352 (Indonesia);

 orcid.org/0000-0002-5493-1041

Neneng Windayani — Department of Chemistry Education, UIN Sunan Gunung Djati, Bandung-40614 (Indonesia);

 orcid.org/0000-0002-7623-9160

Agung Abadi Kiswandono — Department Chemistry, Lampung University, Bandar Lampung-35141 (Indonesia);

 orcid.org/0000-0002-8145-1959

Heri Satria — Department Chemistry, Lampung University, Bandar Lampung-35141 (Indonesia);

 orcid.org/0000-0003-0779-5830

Author Contributions

Conceptualization, S. M. S. and N. N.; Methodology, S. M. S., N. N., and N.W.; Software, S. M. S.; Validation, S. M. S and N. N.; Formal Analysis, S. M. S. and N. N.; Investigation, S. M. S.; Resources, S. M. S.; Data Curation, S. M. S., N. N. and S. S.; Writing – Original Draft Preparation, S. M. S.; Writing – Review & Editing, S. M. S., N. N., S. S., N. W., A. A. K., and H. S.; Project Administration, S. M. S. N. N., S. S., N. W., A. A. K., and H. S.

Conflicts of Interest

The author(s) declare no conflict of interest.

REFERENCES

- [1] J. J. Izquierdo-Gonzalez, F. Amil-Ruiz, S. Zazzu, R. Sanchez-Lucas, C. A. Fuentes-Almagro, and M. J. Rodriguez-Ortega. (2019). "Proteomic analysis of goat milk kefir: Profiling the fermentation-time dependent protein digestion and identification of potential peptides with

- biological activity". *Food Chemistry*. **295** : 456-465. [10.1016/j.foodchem.2019.05.178](https://doi.org/10.1016/j.foodchem.2019.05.178).
- [2] C. Gokirmakli and Z. B. Guzel-Seydim. (2022). "Water kefir grains vs. milk kefir grains: Physical, microbial and chemical comparison". *Journal of Applied Microbiology*. **132** (6): 4349-4358. [10.1111/jam.15532](https://doi.org/10.1111/jam.15532).
- [3] A. Sindi, M. B. Badsha, B. Nielsen, and G. Unlu. (2020). "Antimicrobial Activity of Six International Artisanal Kefirs Against *Bacillus cereus*, *Listeria monocytogenes*, *Salmonella enterica* serovar Enteritidis, and *Staphylococcus aureus*". *Microorganisms*. **8** (6). [10.3390/microorganisms8060849](https://doi.org/10.3390/microorganisms8060849).
- [4] S. Guha, H. Sharma, G. K. Deshwal, and P. S. Rao. (2021). "A comprehensive review on bioactive peptides derived from milk and milk products of minor dairy species". *Food Production, Processing and Nutrition*. **3** (1). [10.1186/s43014-020-00045-7](https://doi.org/10.1186/s43014-020-00045-7).
- [5] D. Tagliazucchi, S. Martini, and L. Solieri. (2019). "Bioprospecting for Bioactive Peptide Production by Lactic Acid Bacteria Isolated from Fermented Dairy Food". *Fermentation*. **5** (4). [10.3390/fermentation5040096](https://doi.org/10.3390/fermentation5040096).
- [6] T. M. Htay, K. K. Sann, and H. Haini. (2023). "Comparative Study on Phytochemical Screening and Antioxidant Activity of Aqueous Extract from Various Parts of *Bauhinia purpurea*". *Bioactivities*. **1** (1): 24-31. [10.47352/bioactivities.2963-654X.183](https://doi.org/10.47352/bioactivities.2963-654X.183).
- [7] A. Biadala and N. M. Adzahan. (2021). "Storage Stability of Antioxidant in Milk Products Fermented with Selected Kefir Grain Microflora". *Molecules*. **26** (11). [10.3390/molecules26113307](https://doi.org/10.3390/molecules26113307).
- [8] L. L. Pruteanu, D. S. Bailey, A. C. Gradinaru, and L. Jantschi. (2023). "The Biochemistry and Effectiveness of Antioxidants in Food, Fruits, and Marine Algae". *Antioxidants (Basel)*. **12** (4). [10.3390/antiox12040860](https://doi.org/10.3390/antiox12040860).
- [9] M. Azizkhani, P. E. J. Saris, and M. Baniasadi. (2020). "An in-vitro assessment of antifungal and antibacterial activity of cow, camel, ewe, and goat milk kefir and probiotic yogurt". *Journal of Food Measurement and Characterization*. **15** (1): 406-415. [10.1007/s11694-020-00645-4](https://doi.org/10.1007/s11694-020-00645-4).
- [10] M. Taniguchi, R. Aida, K. Saito, A. Ochiai, S. Takesono, E. Saitoh, and T. Tanaka. (2019). "Identification and characterization of multifunctional cationic peptides from traditional Japanese fermented soybean Natto extracts". *Journal of Bioscience and Bioengineering*. **127** (4): 472-478. [10.1016/j.jbiosc.2018.09.016](https://doi.org/10.1016/j.jbiosc.2018.09.016).
- [11] K. F. Chai, A. Y. H. Voo, and W. N. Chen. (2020). "Bioactive peptides from food fermentation: A comprehensive review of their sources, bioactivities, applications, and future development". *Comprehensive Reviews in Food Science and Food Safety*. **19** (6): 3825-3885. [10.1111/1541-4337.12651](https://doi.org/10.1111/1541-4337.12651).
- [12] M. de Lima, R. A. da Silva, M. F. da Silva, P. A. B. da Silva, R. Costa, J. A. C. Teixeira, A. L. F. Porto, and M. T. H. Cavalcanti. (2018). "Brazilian Kefir-Fermented Sheep's Milk, a Source of Antimicrobial and Antioxidant Peptides". *Probiotics and Antimicrobial Proteins*. **10** (3): 446-455. [10.1007/s12602-017-9365-8](https://doi.org/10.1007/s12602-017-9365-8).
- [13] Z. Hanum, C. A. Fitri, and Y. Yurliasni. (2021). "Kefir Susu Kambing dengan Penambahan Ekstrak Etanol Kembang Telang (*Clitoria ternatea*) Berpotensi Kuat sebagai Antioksidan dan Antibakteri". *Jurnal Veteriner*. **22** (3): 406-413. [10.19087/jveteriner.2021.22.3.406](https://doi.org/10.19087/jveteriner.2021.22.3.406).
- [14] C. Zong, X. Zhang, F. Yang, Y. Zhou, N. Chen, Z. Yang, G. Ding, F. Yu, and Y. Tang. (2019). "Biotransformation of a crizotinib intermediate using a mutant alcohol dehydrogenase of *Lactobacillus kefir* coupled with glucose dehydrogenase". *Preparative Biochemistry & Biotechnology*. **49** (6): 578-583. [10.1080/10826068.2019.1591987](https://doi.org/10.1080/10826068.2019.1591987).
- [15] Sulmiyati, N. Said, D. U. Fahrodi, R. Malaka, and F. Maruddin. (2019). "Assessment of the antibacterial activity of goat milk kefir on *Escherichia coli* ATCC 8739

- and *Salmonella enterica* subsp. *enterica* serovar typhimurium ATCC 14028 using a well diffusion method". *IOP Conference Series: Earth and Environmental Science*. **247**. [10.1088/1755-1315/247/1/012051](https://doi.org/10.1088/1755-1315/247/1/012051).
- [16] K. T. Magalhães, G. V. d. M. Pereira, C. R. Campos, G. Dragone, and R. F. Schwan. (2011). "Brazilian kefir: structure, microbial communities and chemical composition". *Brazilian Journal of Microbiology*. **42** (2): 693-702. [10.1590/s1517-83822011000200034](https://doi.org/10.1590/s1517-83822011000200034).
- [17] F. Ajam and H. Koohsari. (2020). "Effect of some fermentation conditions on antibacterial activity of fermented milk by kefir grains". *Journal of Food Processing and Preservation*. **44** (12). [10.1111/jfpp.14913](https://doi.org/10.1111/jfpp.14913).
- [18] A. P. d. Oliveira, G. A. d. Santos, C. S. Nomura, and J. Naozuka. (2019). "Elemental chemical composition of products derived from kefir fermented milk". *Journal of Food Composition and Analysis*. **78** : 86-90. [10.1016/j.jfca.2019.02.005](https://doi.org/10.1016/j.jfca.2019.02.005).
- [19] C. Mahdi, H. Untari, and M. C. Padaga. (2018). "Identification and Characterization of Bioactive Peptides of Fermented Goat Milk as a Sources of Antioxidant as a Therapeutic Natural Product". *IOP Conference Series: Materials Science and Engineering*. **299**. [10.1088/1757-899x/299/1/012014](https://doi.org/10.1088/1757-899x/299/1/012014).
- [20] A. C. Sabbione, A. E. Nardo, M. C. Añón, and A. Scilingo. (2016). "Amaranth peptides with antithrombotic activity released by simulated gastrointestinal digestion". *Journal of Functional Foods*. **20** : 204-214. [10.1016/j.jff.2015.10.015](https://doi.org/10.1016/j.jff.2015.10.015).
- [21] I. Hasmila, H. Natsir, and N. H. Soekamto. (2019). "Phytochemical analysis and antioxidant activity of soursop leaf extract (*Annona muricata* Linn.)". *Journal of Physics: Conference Series*. **1341** (3). [10.1088/1742-6596/1341/3/032027](https://doi.org/10.1088/1742-6596/1341/3/032027).
- [22] S. Adi Prayitno, L. Puji Lestari, D. Retnaningtyas Utami, and N. Salsabila. (2021). "Evaluation of Phytochemicals And Antioxidant Activity (IC50) of Bintaro Fruit Ethanol Extract (*Cerbera odollam* L.)". *Food Science and Technology Journal (Foodscitech)*. 1-7. [10.25139/fst.v4i1.3686](https://doi.org/10.25139/fst.v4i1.3686).
- [23] A. Baschieri and R. Amorati. (2021). "Methods to Determine Chain-Breaking Antioxidant Activity of Nanomaterials beyond DPPH(*). A Review". *Antioxidants (Basel)*. **10** (10). [10.3390/antiox10101551](https://doi.org/10.3390/antiox10101551).
- [24] P. Darvishzadeh, V. Orsat, and J. L. Martinez. (2021). "Process Optimization for Development of a Novel Water Kefir Drink with High Antioxidant Activity and Potential Probiotic Properties from Russian Olive Fruit (*Elaeagnus angustifolia*)". *Food and Bioprocess Technology*. **14** (2): 248-260. [10.1007/s11947-020-02563-1](https://doi.org/10.1007/s11947-020-02563-1).
- [25] S. H. Mariam and A. Hernández. (2021). "Isolation and Characterization of Gram-Negative Bacterial Species from Pasteurized Dairy Products: Potential Risk to Consumer Health". *Journal of Food Quality*. **2021** : 1-10. [10.1155/2021/8876926](https://doi.org/10.1155/2021/8876926).
- [26] X. Pan, F. Chen, T. Wu, H. Tang, and Z. Zhao. (2009). "The acid, bile tolerance and antimicrobial property of *Lactobacillus acidophilus* NIT". *Food Control*. **20** (6): 598-602. [10.1016/j.foodcont.2008.08.019](https://doi.org/10.1016/j.foodcont.2008.08.019).
- [27] H. Sharma, G. D. El Rassi, A. Lathrop, V. B. Dobрева, T. S. Belem, and R. Ramanathan. (2021). "Comparative analysis of metabolites in cow and goat milk yoghurt using GC-MS based untargeted metabolomics". *International Dairy Journal*. **117**. [10.1016/j.idairyj.2021.105016](https://doi.org/10.1016/j.idairyj.2021.105016).
- [28] V. Ganatsios, P. Nigam, S. Plessas, and A. Terpou. (2021). "Kefir as a Functional Beverage Gaining Momentum towards Its Health Promoting Attributes". *Beverages*. **7** (3). [10.3390/beverages7030048](https://doi.org/10.3390/beverages7030048).